



Synthesis of 2-(2-hydroxyaryl)alkenylphosphonium salts from phosphine oxides *via* ring-closing ring-opening approach and their antimicrobial evaluation

Dmitry A. Tatarinov^{a, b, *}, Denis M. Kuznetsov^{a, b}, Alexandra D. Voloshina^a, Anna P. Lyubina^{a, b}, Anastasiya S. Strobrykina^a, Fahima K. Mukhitova^c, Fedor M. Polyancev^a, Vladimir F. Mironov^{a, b}

^a A.E. Arbuzov Institute of Organic and Physical Chemistry, Kazan Scientific Center, Russian Academy of Sciences, Arbuzov Str. 8, Kazan, 420088, Russian Federation

^b Kazan Federal University, Kremlevskaya Str. 18, Kazan, 420008, Russian Federation

^c Kazan Institute of Biochemistry and Biophysics, Russian Academy of Sciences, Lobachevsky Str. 2/31, Kazan, 420111, Russian Federation

ARTICLE INFO

Article history:

Received 8 September 2016

Received in revised form

27 October 2016

Accepted 7 November 2016

Available online 9 November 2016

Keywords:

Phosphonium salts

Grignard reagents

Cyclization

Ring opening

Quasiphosphonium salts

Phosphine oxides

ABSTRACT

A new two-step synthesis of phosphonium salts from phosphine oxides using Grignard reagents is reported. This approach involves a cyclization of Z-dialkyl-(diaryl)[2-(5-chloro-2-hydroxyphenyl)-2-phenylvinyl]phosphine oxides under the action of sulfinyl chloride with the formation of 2,2-dialkyl-(diaryl)-6-chloro-4-phenyl-2H-1,2-benzoxaphosphinin-2-ium chlorides followed by ring opening under the action of organomagnesium compounds. The method was successfully applied to prepare a series of a new phosphonium salts bearing phenolic moiety with a wide range of substituents at phosphorus atom. Synthesized phosphonium salts shows high antibacterial and antifungal *in vitro* activity and low toxicity towards human red blood cell.

© 2016 Elsevier Ltd. All rights reserved.

1. Introduction

Phosphonium salts are an extensively studied class of organophosphorus compounds. They have found application in asymmetric phase-transfer catalysis,^{1–4} as organocatalyst⁵ as well as ionic liquids.^{6,7} The introduction of phosphonium functionality onto either natural or synthetic frameworks may lead to new pharmaceutically attractive compounds with antitumor^{8–12} and antimicrobial activities.^{13–15} Recently the structure, synthesis and practical application of polymeric materials containing a quaternary phosphonium moiety as antimicrobial agents have been observed.¹⁶ In addition, some phosphonium salts are able to the reversible inhibition of cholinesterases^{16,17} and possess a

mitochondrion targeted antioxidant activity.^{18–24}

The reactions of P(III) derivatives with alkyl halides or carbonyl compounds in the presence of hydrogen halide are commonly used for the obtaining of the phosphonium salts.^{13,14} In such a reactions triphenyl- or tributylphosphines or rarely other trialkyl derivatives are usually applied. A few works on synthesis of phosphonium salts based on the reaction of tertiary phosphines with ortho-quinones^{25,26} should also be noted. Recent advances of these synthetic approaches are widely reviewed.^{13,14}

Earlier, we have shown by only one example that the reaction of 2-(5-chloro-2-hydroxyphenyl)-2-phenylvinyl diethylphosphine oxide with excess of bromine leads to the cyclic phosphonium salts.²⁷

Here, we report a two-step synthesis of the functionally substituted phosphonium salts **9**, **10** bearing a hydroxy-group in the position delta to phosphorus from quasiphosphonium salts **2**. This approach has been reported briefly earlier as the conference proceedings without any experimental details.²⁸

* Corresponding author. A.E. Arbuzov Institute of Organic and Physical Chemistry, Kazan Scientific Center, Russian Academy of Sciences, Arbuzov Str. 8, Kazan, 420088, Russian Federation.

E-mail address: datint@iopc.ru (D.A. Tatarinov).